

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

03 OCT 2000

WIPO

PCT

5200/1390

PA 267641

THE UNITED STATES OF AMERICA**TO ALL TO WHOM THESE PRESENTS SHALL COME:****UNITED STATES DEPARTMENT OF COMMERCE****United States Patent and Trademark Office****June 29, 2000**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE UNDER 35 USC 111.**

APPLICATION NUMBER: 60/143,632**FILING DATE: July 14, 1999****PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**

T. Wallace
T. WALLACE

Certifying Officer

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Docket Number		003300-580		Type a plus sign (+) inside this box		+	
INVENTOR(s)/APPLICANT(s)							
LAST NAME		FIRST NAME		MIDDLE INITIAL		RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)	
Lindholm		Leif				Östra Björnvägen 8 S-429 30 Kullavik, Sweden (Swedish Nationality)	
TITLE OF THE INVENTION (280 characters max)							
RECOMBINANT ADENOVIRUS							
CORRESPONDENCE ADDRESS							
Benton S. Duffett, Jr. BURNS, DOANE, SWECKER & MATHIS, L.L.P. P.O. Box 1404 Alexandria,							
STATE		Virginia		ZIP CODE		22313-1404	
COUNTRY		United States of America					
ENCLOSED APPLICATION PARTS (check all that apply)							
<input checked="" type="checkbox"/> Specification		Number of Pages		20		<input type="checkbox"/> Small Entity Statement	
<input checked="" type="checkbox"/> Drawing(s)		Number of Sheets		5		<input checked="" type="checkbox"/> Other (specify) Claims 1-23 Abstract of the Disclosure Sequence Listing (12 pages)	
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)							
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees				PROVISIONAL FILING FEE AMOUNT(S)		\$ <input type="checkbox"/> \$75.00	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any deficiency in filing fees or credit any overpayment to Deposit Account Number 02-4800. This paper is submitted in triplicate.						\$ <input checked="" type="checkbox"/> \$150.00	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

- ☒ No.
☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE

Robin L. Teskin

Date July 14, 1999

TYPED or PRINTED NAME Robin L. Teskin for Benton S. Duffett, Jr.

Registration No. 35,030
(if appropriate)

- ☐ Additional inventors are being named on separately numbered sheets attached hereto

RECOMBINANT ADENOVIRUSField of the invention

5 The present invention relates to new recombinant
adenovirus with changed tropism. More particularly the
recombinant adenovirus has been constructed by removing
the native knob structure and replacing it with a new
cell binding ligand and an external trimerisation motif.
10 The invention also relates to the new adenovirus for
treatment of human diseases. Also included is a method
for rescuing of recombinant adenovirus fibers into the
adenovirus genome.

Background of the invention.

15 Clinical gene therapy was introduced in 1989. The aim at
that time was to correct a gene defect in the immune
system through the in vitro introduction of a healthy
gene into the defect cells of the patient and transfusion
20 of the treated cells back to the patient. Since that
time, the possible indications for gene therapy have
increased dramatically. Today, ten years after its
introduction, the use of gene therapy to treat e.g.
diseases of the blood vessels, cancer, inflammatory
25 diseases and infectious diseases such as HIV can be
envisaged.

At present, however, gene therapy is not a useful method
in human medicine. One main reason is that gene therapy
30 demands the packaging of the genes to be delivered into
gene-carriers, or vectors, which can be injected into

60143632-071109

patients and which will target the genes only to the intended cells. Such vectors have so far not been available.

- 5 Adenoviruses (Ad) are DNA viruses without an envelope, shaped as regular icosahedrons with a diameter of 60-85 nm. Cell-binding takes place through fiber proteins, anchored to the virion at the corners of the icosahedron.

- 10 The fiber protein is not necessary for assembly and release of intact virions. Assembly of virions take place in the nucleus of infected cells.

- 15 The fiber protein, which is a homotrimer of a fiber polypeptide, contains three functionally different parts: an N-terminal tail anchoring the fiber non-covalently to the penton base in the virion and which furthermore contains the nuclear-localization signal; an approximate 15 amino acid fiber shaft motif which is repeated six times in Ad3 and 22 times in Ad2 and Ad5 (Chrobozek J, 20 Ruigrok RWH and Cusack S: Adenovirus Fiber, *Current Topics in Microbiology and Immunology*, 1995, p 163-200); and a C-terminal globular domain, the knob, which contains the ligand which binds to the cellular Ad-receptor (See review in in the previous ref.). Each shaft 25 repeat has two three-amino acid regions which form β -sheets and two amino acid regions which constitute the turns of the native extended fiber shaft. The crystal structure of the trimerised, cell-binding domain has been determined and shows a unique topology different from 30 other anti-parallel β -sandwiches (Di Xia, Henry LJ, Gerard RD and Deisenhofer J: Crystal structure of the receptor-binding domain of adenovirus type 5 fiber protein at 1.7 Å resolution; *Structure* 2: 1259-1270;

It seems possible that the fiber can tolerate structural modifications as long as the ability to bind to the penton base and to be transported to the nucleus is

10 retained. Some attempts at modifying the Ad fiber in
order to change the binding properties of the virus have
been made. A short peptide ligand has been added C-
terminally of the knob (Michael SI, Hoy JS, Curie DT and
Engles JT; Addition of a short peptide ligand to the
15 adenovirus fiber protein. *Gene Therapy* 2: 660-8, 1995.)
and an octapeptide has been introduced into one of the
knob "loops". By introducing the FLAG tetra-amino acid
motif into the Ad penton, it has been shown possible to
target Ad to cells normally not infected by Ad. This was
20 done by targeting with bi-specific antibodies where one
specificity was directed against the FLAG peptide and the
other against the new target cell (Wickham TJ, Segal DM,
Roelvink PW, Carrion ME, Lizonova A, Lee GM and Kovesdi
I: Targeted Adenovirus Gene Transfer to Endothelial and
25 Smooth Muscle Cells by Using Bispecific Antibodies. *J.*
Virol., 70: 6831-6838, 1996.). It would therefore seem
possible to target Ad to a broad range of human cells
which would be very useful for the purpose of human gene
therapy. For these reasons and for the reason that Ad
30 have been used extensively for gene therapeutic
applications (Trapnell BC and Gorziglia: Gene therapy
using adenoviral vectors, *Current Opinion in*
Biotechnology, 5: 617-625, 1994.) a method has now been

developed to create recombinant re-targeted Ad-virus which can be useful for human gene therapy.

- Accordingly it is an object of the present invention to
5 provide a recombinant adenovirus with changed tropism.

Another object of the invention is the recombinant adenovirus for treatment of human diseases.

- 10 A further object of the invention is a method for rescuing of recombinant adenovirus fibers into the adenovirus genome.

Summary of the invention

- 15 The objects of the invention are obtained by the recombinant adenovirus and the method for rescuing the virus fibers as claimed in the claims.
- 20 According to the invention there is provided a recombinant adenovirus with changed tropism. The adenovirus is characterized in that the native pentone fibre, which comprises a fibre tail, a fibre shaft and a fibre knob including a trimerisation motif, has been
25 changed in that the native knob containing the cell binding structure and the native trimerisation motif has been removed and a new cellbinding ligand and an external trimerisation motif have been introduced into the virus fiber.
- 30 The structural modification has been performed by DNA technology at the gene level or by chemical or immunological means at the virus level.
- 35 According to another aspect of the invention adenovirus, as identified above, is used for the treatment of human diseases, either in vivo or by in vitro methods.

THE INVENTION IS DESCRIBED IN MORE DETAIL IN THE FOLLOWING

A further aspect of the invention is a method for rescuing of recombinant adenovirus fibers into the adenovirus genome comprising the following steps:

5

- a) subcloning of a 9kb fragment (from S_{pe}I to end of genome),
- b) further subcloning of a 3kb fragment between S_{ac}I and K_pnI,

- 10 c) deletion of the fiber gene between N_{de}I and M_{un}I and replacing the missing sequence with SEQ ID NO: 13 in the Sequence listing containing an X_{ho}I site;
- d) ligation of recombinant fiber between N_{de}I and X_{ho}I of construct under c) above;
- 15 e) re-introduction of construct under d) above into the 9 kb fragment cut with N_{he}I using homologous recombination in *E. coli*;
- f) isolation of the recombinant 9 kb fragment under e) and re-creation of the adenovirus genome by joining 9 kb
- 20 fragment to the 27 kb fragment from the beginning of the genome to the S_{pe}I site by Cosmid cloning.

Detailed description of the invention

25 Figure legends

Fig. 1: Summary of modifications to native fiber carried out in the invention.

30 Fig. 2: Recombinant adenovirus fibers.

Fig. 3: Method for rescuing of recombinant fiber genes into the Ad genome.

35 Fig. 4a: Recombinant fibers rescued into Ad genomes which are capable of giving CPE/plaques on transfected cells and in secondary cultures.

664143632-071499

vectors for cancer gene therapy, *Gene Therapy* 3: 557-559). This may theoretically result in an increase of the chosen effector mechanism over that obtainable with replication incompetent vectors. Furthermore, infectious virus may contribute to an anti tumor effect by cytopathogenic effects in infected cells as well as by evoking an anti viral immune response which may harm infected cells.

10 Construction, expression and evaluation of recombinant fibers

The aim has been to develop a universal method for the construction of functional Ad fibers with changed binding-specificity to make possible the construction of re-targeted Ad.

20 The adenovirus fiber peptide carries several biological functions which are necessary to retain in order to produce active virus particles. The following fiber features are deemed to be of key importance in the construction of functional recombinant fiber peptides:

- The ability to form pentameric homotrimers. This function is carried by the N-terminal amino acid sequence of the wild type fiber knob and is necessary for the fiber to be able to bind to penton base and to form the functional cell binding knob.
- The ability to bind to penton base to form penton capsomeres. This function is carried by the wild type fiber tail.
- The ability to express a cell-binding ligand allowing for attachment to target cells. This function is carried by the wild type fiber knob.

60143632-071499

- Since adenovirus is assembled in the nucleus of infected cells, the ability to be transported into the nucleus of infected cells is vital to virus formation. The nuclear localization signal is mainly, but perhaps not exclusively, carried by the wild type fiber tail.

In the first stage recombinant fibers are constructed and evaluated in vitro after cell-free expression in a

-
- coupled transcription/translation system. Analysis by SDS-PAGE and autoradiography is performed to reveal the presence of an open reading frame and give information on the size of the translated product. In the next stage recombinant fibers are cloned into Baculovirus and expressed in insect cells allowing for functional studies of the fibers. Such studies include ability to form trimers as evaluated by immunostaining with monoclonal antibody 2A6.36 which has been shown to react only with trimerised fibers (Shin Hong J and Engler JA: The amino terminus of the adenovirus fiber protein encodes the nuclear localization signal, *Virology* 185: 758-767, 1991), expression of functional ligand as evidenced by ability to bind to cells expressing the corresponding receptor and ability to bind to penton-base either in solution or on virions.
- Recombinant fibers are constructed using methodology based on PCR (Clackson T, Güssow D and Jones PT: General application of PCR to gene cloning and manipulation, in PCR, A Practical Approach, Eds McPherson MJ, Quirke P and Taylor GR, IRL Press, Oxford, p 187, 1992), e.g. PCR-ligation-PCR (Alvaro Ali S, Steinkasserer A: PCR-ligation-PCR Mutagenesis: A Protocol for Creating Gene Fusions and Mutations, *BioTechniques* 18: 746-750, 1995)

60143632-071499

and splicing by overlap extension (SOE) (Horton RM and Pease LR: Recombination and mutagenesis of DNA sequences using PCR, in McPherson MJ (ed), Directed Mutagenesis, IRL Press 1991, p 217.). Cloning is performed according

- 5 to standard methods. Recombinant fibers are sequenced using Perkin Elmer ABI Prism and are expressed in mammalian cells and in insect cells and stained with monoclonal antibodies specific for fiber tail, trimeric fiber and the new cell-binding ligand. The following
- 10 parameters are evaluated after immunostaining:
- trimerisation
 - nuclear transportation
 - expression of the new cell-binding ligand.

- 15 Finally, recombinant fibers are rescued into the Ad genome by a newly invented procedure described below and recombinant virus particles are produced.

- The invention will be further illustrated with the
- 20 following non-limiting examples:

Example 1:

- Fiber peptides are made where the knob is replaced with
- 25 an external trimerisation motif which is introduced after the TLWT motif which ends the shaft portion of the fiber. The purpose behind the introduction of an external trimerisation motif is two-fold: a) to remove the knob containing the native trimerisation signal but also the
- 30 cell-binding part of the fiber, and b) simultaneously to supply the necessary trimerisation signal. Two different amino acid motifs have been used, i.e. the 36 amino acid "Neck Region Peptide" = NRP, SEQ ID NO: 1 in Sequence

- listing) from human "Lung Surfactant Protein D" (. Hoppe H-J, Barlow PN and Reid KBM: A parallel three stranded - helical bundle at the nucleation site of collagen triple-helix formation. *FEBS Letters* 344: 191-195 (1994).) and a
- 5 31 aa "Zipper" motif where the leucine residues on positions 1 and 4 have been replaced with isoleucine residues = pII (SEQ ID NO: 2 in Sequence listing) (Harbury PB, Tao Zhang, Kim PS and Alber T: A Switch
-
- 10 Between Two-, Three-, and Four-Stranded Coiled Coils in GCN4 Leucine Zipper Mutants. *Science* 262: 1401-1407, 1993.).

The DNA sequences for these trimerisation motifs are synthesized, cloned and sequenced in the project.

- 15 To replace the cellbinding function of the knob a new cellbinding ligand is introduced into the fiber in addition to the external trimerisation amino acid motif.
- 20 To augment the efficiency of nuclear transportation of recombinant fibers an external nuclear localisation sequence is added to the fiber in some cases.
- 25 In further embodiments the fiber in addition contains sequences which increase the survival of the fiber in the cytosol of infected cells, thereby enhancing transportation into the nucleus and virus assembly. Such sequences are e.g. sequences that are present in the wild type knob or in SEQ ID NO: 10 - 12.
- 30

The following types of fibers are constructed using the methods mentioned above (see Fig 2). The sequence of the

SEQUENCE OF THE VECTORS AND THE CONSTRUCTION OF THE FIBERS

11
wild type fiber is shown in the sequence listing as SEQ
ID NO 14.

Type A

5 where the trimerisation motif is fused to the fiber gene
downstream of the fiber shaft after the TLWT motif which
constitutes the four first amino acids of the fiber knob
or downstream of the second turn (Turn b) of any shaft
repeat, the remaining shaft repeats having been removed.

10 The new cellbinding ligand is introduced downstream of
the trimerisation signal with an amino acid linker motif
being added between the trimerisation signal and the
cellbinding ligand.

15 Type B

similar to type A but with a linker motif introduced
immediately upstream of the trimerisation signal.

Type C

20 where the trimerisation motif is introduced after the
first shaft repeat and in turn followed the shaft repeats
17 through 21. The new cellbinding ligand is introduced
downstream of the trimerisation signal with an amino acid
linker motif being added between the trimerisation signal
25 and the cellbinding ligand.

Type D

where the cellbinding ligand is introduced between the
restriction sites NheI and HpaI in the fiber shaft, with
30 an amino acid linker being added both upstream and
downstream of the ligand.

60143633-07449

Type D/Δ

This is a variant of Type D where the fiber shaft downstream of the cellbinding ligand in Type D was removed. Type D and (D/Δ) are constructed with the normal
 5 knob and with the knob being replaced with an external trimerisation signal as in Types A and B.

Type E

which are similar to Type A but with part of the knob
 10 being retained immediately upstream of the external trimerisation motif.

The following amino acid motifs are used as linkers in the above fiber constructs:

15

- SEQ ID NO: 3, derived from Pseudomonas exotoxin
- SEQ ID NO: 4, derived from tissue prothrombin activator
- SEQ ID NO: 5, derived from the hinge region of mouse immunoglobulin

20

- SEQ ID NO: 6, derived from Staphylococcal protein A
- SEQ ID NO: 7, derived from the hinge region of human IgG3
- SEQ ID NO: 8, derived from shaft repeat no 17 of human Ad5

25

Recombinant fibers are cloned into Baculovirus and expressed in Sf9 cells and/or cloned into the vector pSecTag and expressed in COS cells as secreted proteins. Expression is monitored by immunostaining with monoclonal
 30 antibodies 4D2.5 (anti Ad5 fiber) and 2A6.36 (anti trimerised Ad5 fiber). Expression and trimerisation is

654720-2296709

obvious in all recombinant fibers irrespective of length and trimerisation motif.

The various fibers which have been constructed and shown to be able to form trimers and express the new cell binding ligand are shown in Table 1. The results show that the invented technology is capable of generating trimerising fibers which express a new cellbinding ligand. It should therefore be possible to make functional virus with such fibers.

Table I. Results from immunostaining of different recombinant fibers

	Fiber	Detecting antibody				
		A22	2A6	a-EGF	a-Ig	a-Id
15	Type A					
	A1 RGD	+	+			
	A1 EGF	+	+	+		
20	A1 G250 HK	+	+		+	+
	A1 G250 KH	+	+		+	+
	A1 G250 KHJCH2	+	+		+	+
	A1 VaLVβCβ	+	+			
25	A7 RGD	+	+			
	A7 EGF	+	+	+		
	A7 G250 HK	+	+		+	+
	A7 G250 KH	+	+		+	+
	A7 G250 KHJCH2	+	+		+	+
30	A7 VaLVβCβ	+	+			
	A7 IgG3 EGF	+	+	+		
	A7 (Gly4Ser)4 G250VKVH	+	+		+	+
35	A22 EGF	+	+	+		
	A22 RGD	+	+			
	Type B					
40	B (Gly4Ser)4 RGD	+	+			
	Type C					
	C IgG3 EGF	+	+	+		

C (Gly4Ser)4-
G250VKVH

+ + + +

Type D

5 N/D EGF

+ + +

N/D G250 HKCKy

+ + + + +

F2/D EGF

+ + +

F3/D EGF

+ + +

10 Type D/A

F2 D/A G250 HKCK

+ + +

~~F2 D/A G250 HKCKy~~

~~+ + +~~

F2 D/A EGF

+ + +

F3 D/A EGF

+ + +

15

Abbreviations used in Table 1:

2A6: antibody against trimerized fiber

4D2: antibody against fiber

a-EGF: antibody against epidermal growth factor

20 a-Id: anti idiotypic antibody specific for G250

a-Ig: antibody against mouse immunoglobulin

C β : Constant domain from β chain of T cell receptor
against MAGE1/HLA A1. SEQ ID NO: 11.

CH2: immunoglobulin heavy chain constant domain 2

25 EGF: epidermal growth factor

G250: monoclonal antibody specific for renal carcinoma

H: heavy chain variable sequence from G250 (SEQ ID NO:
15)30 IgG3: amino acid linker derived from hinge region of
human IgG3, SEQ ID NO: 7

J: immunoglobulin joining chain sequence

K: light chain variable sequence from monoclonal antibody
G250 (SEQ ID NO: 16)RGD: The amino acid sequence arginine-glycine-aspartic
acid35 V α : Variable domain from α chain of T cell receptor
against MAGE1/HLA A1. SEQ ID NO: 10V β : Variable domain from β chain of T cell receptor
against MAGE1/HLA A1. SEQ ID NO: 12

60143632 071499

Example 2:

Nuclear localization of recombinant fibers (Tables 2 and 3)

- 5 Nuclear localization is assessed by immunostaining of fibers in Sf9 cells 24 hours after infection with the relevant Baculovirus clone. Some results are shown in Table 2 below. It is clear from these experiments that some recombinant fibers show a grossly impaired nuclear
- 10 localization in Sf9 cells despite the presence of the nuclear addressing signal in the fiber tail.

15

Table 2
Nuclear localization of native and selected recombinant fibers in Sf9 cells

Fiber	% of fiber-expressing Sf9 cells showing nuclear localization after infection
20 Wild type	100
N/D EGF	100
A RGD	App. 50
A7 RGD	App. 100
25 A7 EGF	App. 100
A7 scTCR	App. 50
A7 G250 scFvs	0

- 30 Recombinant and native fibers have also been expressed in COS cells, targeted for expression in the cytosol after cloning into the vector pcDNA 3.1. In this case it was expected that the fibers would be detected in the nucleus, due to the presence of the native nuclear
- 35 localization signal in the fiber tail. However, nuclear localization has so far only been detected in the wild type fiber and in fibers with single-chain T-cell

654720-2294105

receptors, i.e. the fibers which have produced the most efficient virus (see below).

Since nuclear localization of fibers are crucial to virus assembly, an attempt is made to improve the efficiency of nuclear addressing by adding an external nuclear localization signal (NLS), in this case the SV40 large T-antigen NLS having the amino acid sequence SEQ ID NO: 9

(Fisher-Fantuzzi L and Vesco C: Cell-Dependent Efficiency of Reiterated Nuclear Signals in a Mutant Simian Virus 40 Oncoprotein Targeted to the Nucleus. *Mol Cell Biol*, 8:5495-5503, 1988). The external NLS sequence is added immediately up-stream of the RGD motif. It is found that the presence of the external NLS dramatically improved the nuclear localization in the cases where it has been investigated. In fact, as mentioned above the fiber constructs lacking the external NLS were undetectable in the transfected cells (Table 3).

20

Table 3

Nuclear localization of native and selected recombinant fibers in COS cells after targeting for expression in the cytosol

25

Fiber	Nuclear localization
Wild type	+
A ValV β C β	+
A ValV β C β CK	+
A RGD	-
A NLS RGD	+
A7 RGD	-
A7 NLS RGD	+
A22 RGD	-

35

For abbreviations, see Table 1

The evidence given above support the hypothesis that recombinant fibers are poorly transported into the nucleus despite the presence of the intact tail region (see also below) and that this may possibly be corrected by the incorporation of an external NLS in the fiber construct.

Example 3:

10 METHOD FOR RESCUING OF RECOMBINANT FIBERS INTO VIRIONS

The wild type fiber in the Ad genome is exchanged for recombinant fibers by the following method (see Fig 3).

The plasmid pTG3602 (Chartier C, Degryse E, Gantzer M, Dieterlé A, Pavirani A and Mehtali M: Efficient generation of Recombinant Adenovirus Vectors by Homologous Recombination in *Escherichia Coli*, *J Virol*, 70: 4805-4810, 1996) containing the entire Ad5 genome as a PacI-PacI fragment is used as starting material. The approximate 9kb fragment of the genome between SpeI and PacI and containing the wild type fiber gene is cloned separately in pBluescript. From this fragment an approximate 3kb fragment between SacI and KpnI is further subcloned. A deletion of the native fiber gene with the exception of the N-terminal nucleotides upstream of the NdeI site of the fiber, between the NdeI site and the MunI site, which begins at base 38 after the stop codon of the fiber, is created in the 3kb fragment. The deleted sequence is replaced with SEQ ID NO: 13 which restores the NdeI and MunI sites and the wild type genome sequence between the fiber stop codon and the MunI site. In addition the added sequence, SEQ ID NO: 13, contains an XhoI site allowing for ligation of recombinant fibers

004420-2294102

into the fiber-deleted 3kb fragment (the 3 kb fiber shuttle) between NdeI and XhoI.

The 3 kb fiber shuttle with recombinant fiber is re-
 5 introduced into the 9 kb fragment cut with NheI using
 homologous recombination in E.coli (see ref. in previous
 passage). The resulting recombinant 9 kb fragment is
 finally excised from the vector with SpeI and PaeI and
 joined to the isolated 27 kb fragment by Cosmid cloning.

10

The presence of an insert and the expected properties is
 verified in all cosmid clones by PCR. Cosmid clones are
 also restricted with Hind III and the presence of
 restriction fragments of the expected size verified on
 15 gels.

Recombinant Ad genomes are isolated after restriction
 with Pac I and used to transfect suitable cells. The
 occurrence of plaques is determined by microscopic
 20 inspection of the transfected cell cultures.

Supernatants are harvested from primarily transfected
 cultures and used to infect secondary cultures. The
 occurrence of cytopathogenic effects and plaques are
 25 monitored by microscopy.

The particular fiber constructs that have been
 successfully rescued into virus are shown in figure 4a
 and 4b.

00143301433

Conclusion:

For gene therapy to be useful for treatment of human
5 diseases there is a need for injectable vectors with
ability to target specific cells or a specific tissue
(Miller N and Vile R: Targeted vectors for gene therapy.
FASEB J, 9: 190-199, 1995).

10 The present invention describes methods whereby knobless,
trimerisation-competent fibers with new cellbinding
ligands can be created and rescued into virus and have
identified locations within the fiber-shaft which
15 tolerates inserts of foreign ligands. The importance of
intracellular trafficking of recombinant fibers has also
been identified. Recombinant virus made using the
invented technology should be highly useful in human
medicine. Virtually unlimited opportunities for targeted
20 gene-therapy may be developed by the combination of the
technology described here and the identification of cell-
binding ligands by phage-display.

So far trimerisation-competent fibers with a human scTCR
have been and rescued into functional virus. Since single
25 chain antibodies are large and highly complex peptides it
seems highly likely that also other scAbs and cell-
binding ligands, e.g. peptides identified from peptide
libraries by means of phage-display, could be
incorporated into Ad-fibers and rescued into virus using
30 the same technology.

There are many ways in which Ad, made re-targeted by the
present invention, may be applied to human gene therapy.

THE PRESENT INVENTION IS NOT LIMITED BY THE ABOVE DISCUSSION.

6014332-071499

In the case of tumor diseases, the following options exist:

I. Use of vectors to introduce transgenes into tumors,
5 such as

- anti onco genes
- "suicide" genes
- genes for immune modulatory substances or tumor
antigens

10 • genes for anti angiogenetic factors

II. Use of infectious virus. This has the added value
over the use of non replicating vectors that virus can
spread from cell to cell within a tumor, thereby
15 multiplying the initial hit on the tumor. Tumor cell
destruction may occur not only by the cell-destroying
mechanism engineered into the vector but also by the cell
destruction which is associated with the virus infection
per se and by the attack of the body's immune response on
20 the virus infected cells. This principle has already been
tested in man through the direct intra-tumoral injection
of an adenovirus which has been made gene manipulated to
replicate only in p53 mutant tumor cells. The experience
from these limited trials on large "head-and-neck" tumors
25 are partially encouraging with a complete regress of 2/11
treated tumors which are otherwise resistant to any form
of known treatment.

65413632-071449

Claims

1. Recombinant adenovirus with changed tropism, characterized in that the native pentone fibre, comprising a fibre tail, a fibre shaft and a fibre knob including a trimerisation motif, has been changed in that the native knob containing the cell binding structure and the native trimerisation motif has been removed and a new cellbinding ligand and an external trimerisation motif have been introduced into the virus fiber.

10

2. Adenovirus according to claim 1, chracterized in that said structural modification has been performed by DNA technology at the gene level or by chemical or immunological means at the virus level.

15

3. Adenovirus according to claim 1 which is either replication competent or replication in-competent.

4. Adenovirus according to claim 1, characterized in that the new cellbinding ligand has been introduced into the fiber shaft.

20

5. Adenovirus according to claim 1, characterized in that the new cell binding ligand has been introduced downstream of the fiber shaft repeats.

25

6. Adenovirus according to claim 4 characterized in that the new cellbinding ligand has been introduced between the restriction sites NheI and HpaI in the fiber shaft.

30

7. Adenovirus according to claim 4, characterized in that amino acid linkers have been introduced upstream and downstream of the cellbinding ligand.

35

60143622-074499

8. Adenovirus according to claim 4, characterized in that the shaft repeats downstream of the restriction site Hpa1 have been removed.

- 5 9. Adenovirus according to claim 1, characterized in that an amino acid linker --- has been added between the fiber shaft and the trimerisation motif and/or between the trimerisation motif and the cellbinding ligand as a linker.

10

10. Adenovirus according to claim 9, characterized in that the amino acid linker motif is any of the following: SEQ ID NO: 3, derived from Pseudomonas exotoxin; SEQ ID NO: 4, derived from tissue prothrombin activator; SEQ ID NO: 5, derived from the hinge region of mouse immunoglobulin; SEQ ID NO: 6, derived from Staphylococcal protein A; SEQ ID NO: 7, derived from the hinge region of human IgG3; SEQ ID NO: 8, derived from shaft repeat 17 of human Ad5.

20

11. Adenovirus according to any of the claims 1 - 10, characterized in that the new cellbinding ligand is any cellbinding peptide.

25

12. Adenovirus according to claim 11, characterized in that the cell binding ligand is a monoclonal antibody or a fragment thereof whether as a single chain fragment or Fab, a T cell receptor or a fragment thereof, an integrin binding peptide such as RGD or a growth factor such as

30 Epidermal Growth Factor.

13. Adenovirus according to claim 12, containing any of the sequences SEQ ID NO: 10 - 12.

35

14. Adenovirus according to claim 12, characterized in that the single chain fragment is a single chain fragment of the monoclonal antibody G250 with heavy chain variable

50143632-071490

region with SEQ ID NO: 15 and light chain variable region with SEQ ID NO: 16.

15. Adenovirus according to claim 1 characterized in
5 that the external trimerisation motif is an α -helical coiled coil motif, or any other peptide capable of rendering functionally trimerised fibers.

16. Adenovirus according to claim 15, characterized in
10 that the external trimerisation motif is the neck region peptide of human lung surfactant protein D, SEQ ID NO: 1 or a 31 aa "Zipper" motif where the leucine residues on positions 1 and 4 have been replaced with isoleucine residues, SEQ ID NO: 2.

- 15 17. Adenovirus according to any of the preceding claims characterized in that an external nuclear localisation signal (NLS) has been introduced in the fiber.

- 20 18. Adenovirus according to claim 17, characterized in that the NLS is the SV40 large-T antigen NLS.

19. Adenovirus according to any of the preceding claims characterized in that the fiber in addition contains
25 sequences which increase the survival of the fiber in the cytosol of infected cells, thereby enhancing transportation into the nucleus and virus assembly.

20. Adenovirus according to claim 19, characterized in
30 that the sequences are present in the wild type knob.

21. Adenovirus according to claim 20, characterized in that the sequences are present in SEQ ID NO: 10 - 12.

- 35 22. Adenovirus according to claims 1 - 21 for the treatment of human diseases, either in vivo or by in vitro methods.

661120-EEETH09

23. A method for rescuing of recombinant adenovirus fibers into the adenovirus genome comprising the following steps:

5

- a) subcloning of a 9kb fragment (from SpeI to end of genome),
- b) further subcloning of a 3kb fragment between SacI and KpnI,

- 10 c) deletion of the fibergene between NdeI and MunI and replacing the missing sequence with the sequence SEQ ID NO: 13 containing an XhoI site;
- d) ligation of recombinant fiber between NdeI and XhoI of construct under c) above;
- 15 e) re-introduction of construct under d) above into the 9 kb fragment cut with NheI using homologous recombination in *E. coli*;
- f) isolation of the recombinant 9 kb fragment under e) and re-creation of the adenovirus genome by joining 9 kb
- 20 fragment to the 27 kb fragment from the beginning of the genome to the SpeI site by Cosmid cloning.

661120-22934109

Abstract

- Recombinant adenovirus with changed tropism. In the adenovirus the native pentone fibre, comprising a fibre tail, a fibre shaft and a fibre knob including a
- 5 trimerisation motif, has been changed in that the native knob containing the cell binding structure and the native trimerisation motif has been removed and a new cellbinding ligand and an external trimerisation motif have been introduced into the virus fiber. Further, the
- 10 invention relates to the recombinant adenovirus for the treatment of human diseases, either in vivo or by in vitro methods and also to a method for rescuing of recombinant adenovirus fibers into the adenovirus genome.

664720-2292409

Sequence listing

<110> Got-A-Gene AB
 <120> Recombinant adenovirus
 <130> 2998645
 <160> 16
 <170> MS Word 97

<210> 1
 <211> 36
 <212> PRT
 <213> Homo sapiens
 <301> Hoppe HJ, Barlow PN, Reid KBM
 <302> A parallel three stranded a-helical bundle at the nucleation site of collagen triple-helix formation
 <303> FEBS Letters
 <304> 344
 <306> 191-195
 <307> 1994
 <400> 1

Pro Asp Val Ala Ser Leu Arg Gln Gln Val Glu Asp Leu Gln Gly
 1 5 10 15

Gln Val Gln His Ley Gln Ala Ala Phe Ser Gln Tyr Lys Lys Val
 20 25 30

Glu Leu Phe Pro Asn Gly
 35

664120-2E9E4109

<210> 2
 <211> 31
 <212> PRT
 <213> Homo sapiens
 <301> Harbury PB, Zhang T, Kim PS, Albert T
 <302> A switch between two-, three-, and four-stranded coiled coils in GCN4
 leucine zipper mutants

<303> Science
 <304> 262
 <306> 1401-1407
 <307> 1993-11-26
 <400> 2

Met Lys Gln Ile Gly Asp Lys Ile Glu Glu Ile Leu Ser Lys Ile Tyr His
 1 5 10 15

Ile Glu Asn Gly Ile Ala Arg Ile Lys Lys Leu Ile Gly Glu
 20 25 30

<210> 3
 <211> 6
 <212> PRT
 <213> Pseudomonas aeruginosa
 <301> Brinkmann U, Buchner J, Pastan I
 <302> Independent domain folding of Pseudomonas exotoxin and single
 chain immunotoxins: Influence of interdomain connections
 <303> Proc Natl Acad Sci US
 <304> 89

667120-2294109

<306> 3075-3079
 <307> 1992
 <400> 3
 Ala Ser Gly Gly Pro Glu
 1 5

<210> 4
 <211> 7
 <212> PRT
 <213> Homo sapiens
 <301> Brinkmann U, Buchner J, Pastan I
 <302> Independent domain folding of Pseudomonas exotoxin and single
 chain immunotoxins: Influence of interdomain connections
 <303> Proc Natl Acad Sci US
 <304> 89
 <306> 3075-3079
 <307> 1992
 <400> Ala Ser Glu Gly Asn Ser Asp
 1 5

<210> 5
 <211> 8
 <212> PRT
 <213> Mus musculus

654720-2994109

<301> Brinkmann U, Buchner J, Pastan I
 <302> Independent domain folding of Pseudomonas exotoxin and single
 chain immunotoxins: Influence of interdomain connections
 <303> Proc Natl Acad Sci US
 <304> 89
 <306> 3075-3079

<307> 1992

<400> 5

Ala Ser Thr Pro Glu Pro Asp Pro
 1 5

<210> 6

<211> 13

<212> PRT

<213> Staphylococcus aureus

<400> 6

Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Ser Asp
 1 5 10

<210> 7

<211> 11

<212> PRT

<213> Homo sapiens

664420-2294109

<301> Dangi JL, Wensel TG, Morrison SL, Streyer L, Herzenberg LA and Oi
T

<302> Segmental flexibility and complement fixation of genetically
engineered chimeric human, rabbit and mouse antibodies

<303> EMBO Journal

<304> 7

<306> 1989

<307> 1988

<400> 7

Thr Pro Leu Gly Asp Thr Thr His Thr Ser Gly
1 5 10

<210> 8

<211> 11

<212> PRT

<213> Adenovirus type 5

<301> Stouten PFW, Sander C, Ruigrok WH, Cusack S

<302> New triple-helical model for the shaft of the adenovirus fibre

<303> Journal of molecular biology

<304> 226

<306> 1073-1084

<307> 1992

<400> 8

Phe Thr Ala Ser Asn Asn Ser Lys Lys Leu Glu
1 5 10

664720-22964109

9

8

PRT

Simian virus 40

Fisher-Eppluzzi L and Vesco C 8:5495-5503, 1988

Cell-Dependent Efficiency of Reiterated Nuclear Signals in a Mutant Simian Virus 40 Oncoprotein Targeted to the Nucleus

Molecular Cell Biology

8

5495-5503

1992

9

Asp Pro Lys Lys Lys Arg Lys Val
1 5 1

10

119

PRT

- **Homo sapiens**

10

Gln Lys Val Thr Gln Ala Gln Thr Glu Ile Ser Val Val Glu Lys Glu
1 5 10 15

Asp Val Thr Leu Asp Cys Val Tyr Glu Thr Arg Asp Thr Thr Tyr
20 25 30

Tyr Leu Phe Trp Tyr Lys Gln Pro Pro Ser Gly Glu Leu Val Phe Leu Ile
35 40 45

Arg Arg Asn Ser Phe Asp Glu Gln Asn Glu Ile Ser Gly Arg Tyr Ser
 50 55 60 65
 Trp Asn Phe Gln Lys Ser Thr Ser Ser Phe Asn Phe Thr Ile Thr Ala
 70 75 80
 Ser Gln Val Val Asp Ser Ala Val Tyr Phe Cys Ala Leu Gly Gly Val
 85 90 95
 Asn Asn Asn Ala Gly Asn Met Leu Thr Phe Gly Gly Gly Thr Arg
 100 105 110
 Leu Met Val Lys Pro
 115

<210> 11

<211> 133

<212> PRT

<213> Homo sapiens

<400> 11

Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu
 1 5 10 15
 Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thre Leu Val Cys
 20 25 30
 Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Lys Ser Trp Trp
 35 40 45
 Val Asn Gly Lys Glu Val His Ser Gly Val Set Thr Asp Pro Gln Pro
 50 55 60
 Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser
 65 70 75
 Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe
 80 85 90
 Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr
 95 100 105 110
 Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly
 115 120 125
 Arg Ala Asp Ala Ala Ala
 130

<210> 12

<211> 114

<212> PRT

<213> Homo sapiens

<400> 12

```

Asp Ser Gly Val Thr Gln Thr Pro Lys His Leu Ile Thr Ala Thr Gly
 1             5             10             15
Gln Arg Val Thr Leu Arg Cys Ser Pro Arg Ser Gly Asp Leu Ser Val
      20             25             30
Tyr Trp Tyr Gln Gln Ser Leu Asp Gln Gly Leu Gln Phe Leu Ile His
 35             40             45
Tyr Tyr Asn Gly Glu Glu Arg Ala Lys Gly Asn Ile Leu Glu Arg Phe
50             55             60             65
Ser Ala Gln Gln Phe Pro Asp Leu His Ser Glu Leu Asn Leu Ser Ser
      70             75             80
Leu Glu Leu Gly Asp Ser Ala Leu Val Phe Cys Ala Ser Asn Ile Ala
      85             90             95
Gly Gly Ser Tyr Thr Gln Tyr Phe Gly Pro Gly Thr Arg Leu Thr Val
100             105             110
Leu

```

<210> 13

<211> 52

<212> DNA

<213> Artificial sequence

<223> Sequence replacing the fiber gene sequence which was deleted between the NdeI restriction site in the fiber tail and the MunI site which begins at base 38 after the stop codon in the fiber. The sequence restores the NdeI and MunI sites and the wild type genome sequence between the fiber stop codon and the MunI site. In addition the added sequence contains an XhoI site allowing for the ligation of recombinant fibers.

<400> 13

tatgcactcg agtaagaat cgtttggtt atgtttcaac gtgtttattt tc

654120-2294109

<210> 14
 <211> 1746
 <212> DNA
 <213> Human adenovirus type 5
 <221> CDS
 <222> 1-1746

<223> 1-129 Fiber tail
 130-1200 Fiber shaft
 1201-1746 Fiber knob

<400> 14

atg aag cgc gca aga ccg tct gaa gat acc ttc aac ccc gtg tat cca 48
 Met Lys Arg Ala Arg Pro Ser Glu Asp Thr Phe Asn Pro Val Tyr Pro
 1 5 10 15
 tat gac acg gaa acc ggt cct cca act gtg cct ttt ctt act cct ccc 96
 Tyr Asp Thr Glu Thr Gly Pro Pro Thr Val Pro Phe Leu Thr Pro Pro
 20 25 30
 ttt gta tcc ccc aat ggg ttt caa gag agt ccc cct ggg gta ctc tct 144
 Phe Val Ser Pro Asn Gly Phe Gln Glu Ser Pro Pro Gly Val Leu Ser
 35 40 45
 ttg cgc cta tcc gaa cct cta gtt acc tcc aat ggc atg ctt gcg ctc 192
 Leu Arg Leu Ser Glu Pro Leu Val Thr Ser Asn Gly Met Leu Ala Leu
 50 55 60
 aaa atg ggc aac ggc ctc tct ctg gac gag gcc ggc aac ctt acc tcc 240
 Lys Met Gly Asn Gly Leu Ser Leu Asp Glu Ala Gly Asn Leu Thr Ser
 65 70 75 80
 caa aat gta acc act gtg agc cca cct ctc aaa aaa acc aag tca aac 288
 Gln Asn Val Thr Thr Val Ser Pro Pro Leu Lys Lys Thr Lys Ser Asn
 85 90 95
 ata aac ctg gaa ata tct gca ccc ctc aca gtt acc tca gaa gcc cta 336
 Ile Asn Leu Glu Ile Ser Ala Pro Leu Thr Val Thr Ser Glu Ala Leu
 100 105 110
 act gtg gct gcc gcc gca cct cta atg gtc gcg ggc aac aca ctc zcc 384
 Thr Val Ala Ala Ala Ala Pro Leu Met Val Ala Gly Asn Thr Leu Thr
 115 120 125
 atg caa tca cag gcc ccg cta acc gtg cac gac tcc aaa ctt agc att 432
 Met Gln Ser Gln Ala Pro Leu Thr Val His Asp Ser Lys Leu Ser Ile
 130 135 140
 gcc acc caa gga ccc ctc aca gtg tca gaa gga aag cta gcc ctg caa 480
 Ala Thr Gln Gly Pro Leu Thr Val Ser Glu Gly Lys Leu Ala Leu Gln
 145 150 155 160

6541322294109

aca tca ggc ccc ctc acc acc acc gat agc agt acc ctt act atc act 528
 Thr Ser Gly Pro Leu Thr Thr Thr Asp Ser Ser Thr Leu Thr Ile Thr
 165 170 175
 gcc tca ccc cct cta act act gcc act ggt age ttg ggc att gac ttg 576
 Ala Ser Pro Leu Thr Thr Ala Thr Gly Ser Leu Gly Ile Asp Leu
 180 185 190
 aaa gag ccc att tat aca caa aat gga aaa cta gga cta aag tac ggg 624
 Lys Glu Pro Ile Tyr Thr Gln Asn Gly Lys Leu Gly Leu Lys Tyr Gly
 195 200 205
 gct cct ttg cat gta aca gac gac cta aac act ttg acc gta gca act 672
 Ala Pro Leu His Val Thr Asp Asp Leu Asn Thr Leu Thr Val Ala Thr
 210 215 220
 ggt cca ggt gtg act att aat act tcc ttg caa act aaa gtt act 720
 Gly Pro Gly Val Thr Ile Asn Asn Thr Ser Leu Gln Thr Lys Val Thr
 225 230 235 240
 gga gcc ttg ggt ttt gat tca caa ggc aat atg caa ctt aat gta gca 768
 Gly Ala Leu Gly Phe Asp Ser Gln Gly Asn Met Gln Leu Asn Val Ala
 245 250 255
 gga gga cta agg att gat tct caa aac aga cgc ctt ata ctt gat gtt 816
 Gly Gly Leu Arg Ile Asp Ser Gln Asn Arg Arg Leu Ile Leu Asp Val
 260 265 270
 agt tat ccg ttt gat gct caa aac caa cta aat cta aga cta gga cag 864
 Ser Tyr Pro Phe Asp Ala Gln Asn Gln Leu Asn Leu Arg Leu Gly Gln
 275 280 285
 ggc cct ctt ttt ata aac tca gcc cac aac ttg gat att aac tac aac 912
 Gly Pro Leu Phe Ile Asn Ser Ala His Asn Leu Asp Ile Asn Tyr Asn
 290 295 300
 aaa ggc ctt tac ttg ttt aca gct tca aac aat tcc aaa aag ctt gag 960
 Lys Gly Leu Tyr Leu Phe Thr Ala Ser Asn Asn Ser Lys Lys Leu Glu
 305 310 315 320
 gtt aac cta agc act gcc aag ggg ttg atg ttt gac gct aca gcc ata 1008
 Val Asn Leu Ser Thr Ala Lys Gly Leu Met Phe Asp Ala Thr Ala Ile
 325 330 335
 gcc att aat gca gga gat ggg ctt gaa ttt ggt tca cct aat gca cca 1056
 Ala Ile Asn Ala Gly Asp Gly Leu Glu Phe Gly Ser Pro Asn Ala Pro
 340 345 350
 aac aca aat ccc ctc aaa aca aaa att ggc cat ggc cta gaa ttt gat 1104
 Asn Thr Asn Pro Leu Lys Thr Lys Ile Gly His Gly Leu Glu Phe Asp
 355 360 365
 tca aac aag gct atg gtt cct aaa cta gga act ggc ctt agt ttt gac 1152
 Ser Asn Lys Ala Met Val Pro Lys Leu Gly Thr Gly Leu Ser Phe Asp
 370 375 380
 agc aca ggt gcc att aca gta gga aac aac aac aac aag cta act 1200
 Ser Thr Gly Ala Ile Thr Val Gly Asn Lys Asn Asn Asp Lys Leu Thr
 385 390 395 400
 ttg tgg acc aca cca gct cca tct cct aac tgt aga cta aat gca gag 1248
 Leu Trp Thr Thr Pro Ala Pro Ser Pro Asn Cys Arg Leu Asn Ala Glu
 405 410 415

664120-2292109

aaa gat get aaa ctc act ttg gtc tta aca aaa tgt ggc agt caa ala 1296
 Lys Asp Ala Lys Leu Thr Val Leu Thr Lys Cys Gly Ser Gln Ile
 420 425 430
 ctt gct aca gtt tca gtt ttg gct gtt aaa ggc agt ttg gct cca ata 1344
 Leu Ala Thr Val Ser Val Leu Ala Val Lys Gly Ser Leu Ala Pro Ile
 435 440 445
 tct gga aca gtt caa agt gct cat ctt att ata aga ttt gac gaa aat 1392
 Ser Gly Thr Val Gln Ser Ala His Leu Ile Ile Arg Phe Asp Glu Asn
 450 455 460
 gga gtg cta cta aac aat tcc ttc ctg gac cca gaa tat tgg aac ttt 1440
 Gly Val Leu Leu Asn Asn Ser Phe Leu Asp Pro Glu Tyr Trp Asn Phe
 465 470 475 480
 aga nat gga gat ctt act gaa ggc aca gcc tat aca aac ggt gtt gga 1488
 Arg Asn Gly Asp Leu Thr Glu Gly Thr Ala Tyr Thr Asn Gly Val Gly
 485 490 495
 ttt atg cct aac cta tca gct tat cca aaa tct cac ggt aaa act gcc 1536
 Phe Met Pro Asn Leu Ser Ala Tyr Pro Lys Ser His Gly Lys Thr Ala
 500 505 510
 aaa agt aac att gtc agt caa gtt tac tta aac gga gac aaa act aaa 1584
 Lys Ser Asn Ile Val Ser Gln Val Tyr Leu Asn Gly Asp Lys Thr Lys
 515 520 525
 cct gla aca cta acc att aca cta aac ggt aca cag gaa aca gga gac 1632
 Pro Val Thr Leu Thr Ile Thr Leu Asn Gly Thr Gln Glu Thr Gly Asp
 530 535 540
 aca act cca agt gca tac tct atg tca ttt tca tgg gac tgg tct ggc 1680
 Thr Thr Pro Ser Ala Tyr Ser Met Ser Phe Ser Trp Asp Trp Ser Gly
 545 550 555 560
 cac aac tac att aat gaa ata ttt gcc aca tcc tct tac act ttt tca 1728
 His Asn Tyr Ile Asn Glu Ile Phe Ala Thr Ser Ser Tyr Thr Phe Ser
 565 570 575
 tac att gcc caa gaa taa
 Tyr Ile Ala Gln Glu ***

<210> 15
 <211> 120
 <212> PRT
 <213> Mus musculus
 <400> 15

Asp Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Lys Leu Gly Gly
 1 5 10 15

604120-2294109

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30
 Tyr Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Leu Val
 35 40 45
 Ala Ala Ile Asn Ser Asp Gly Gly Ile Thr Tyr Tyr Leu Asp Thr Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75
 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Leu Phe Tyr Cys
 80 85 90 95
 Ala Arg His Arg Ser Gly Tyr Phe Ser Met Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Ser Val Thr Val Ser Ser Gly Ser
 115

<210> 16
 <211> 116
 <212> PRT
 <213> Mus musculus
 <400> 16

Asp Ile Val Met Thr Gln Ser Gln Arg Phe Met Ser Thr Thr Val Gly
 1 5 10 15
 Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asn Val Val Ser Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Met Gln Ser
 65 70 75 80
 Glu Asp Leu Ala Asp Phe Phe Cys Gln Gln Tyr Ser Asn Tyr Pro Trp
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala Asp Ala Ala
 100 105 110
 Pro Thr Val Ser
 115

60143632-071499



AWAPATENT

Handläggs av
Marie-Louise Ebbinghaus

Stockholm
1999-07-05

Referens
2998645

Ansökningsnr.

Härmed intygas, att samtidigt härmed inlämnade diskett rörande sekvenslista i ovannämnda patentansökningsärende har motsvarighet i den i ärendet ingivna sekvenslistan i utskriven form.

FL-E

Ansvarig teknisk medarbetare

66440-22984109


Korrespondensadress:

AWAPATENT AB
Box 45086
104 30 STOCKHOLM


Huvudkontor och styrelsens säte Malmö

telefon 08-440 95 00
telefax 08-440 95 50
epost mail@awapatent.com
Org. nr. 556082-7023


Wild type fiber trimer



Fiber-penton interaction

Mediates
Internalisation of
virus via Integrins
on the cell surfaceCAR Receptor-
fiber interaction
mediates binding
to the cell surface

Recombinant fiber

Cellular receptor
binding to new
ligand on
recombinant fibers

Code

Fiber tail

Fiber shaft

Fiber knob

New trimerisation motif

Linker motif


New cellbinding ligand

Native trimerisation motif


Nuclear localisation signal

Fig. 1

Wild type fiber.



Type A




Type A1. Shaft repeat 1.



Type A7. Shaft repeats 1-7



Type B



Type C



R1 R18-21

Type D



R 1-8

R18-21








Type DΔ. Variant lacking R18-21



Type E. Contains A, B and C sheets of knob.



Code

-  Fiber tail
-  Fiber shaft
-  Fiber knob
-  New trimerisation motif
-  Linker motif
-  New cellbinding ligand
-  Native trimerisation motif

R = Shaft repeat

Linker motifs

ASGGPE = Pseudo exo
 ASEGNSD = TPA
 ASTPEPDP = Ab Hinge, mouse
 AKKLNDQAQPKSD from SpA
 TPLGDTTHTSG = Upper hinge from human IgG3
 (GGGGS)₄

Fig. 2

664120-22924109





Fig. 3

Type A

Time in days for development of
plaques on primarily transfected
cells



Color code




Fig. 4a

6644632-074499

*Time in days for development of
plaques on primarily transfected
cells*

Type B



21



23



No plaques



12




11



No plaques

Type C



Uncertain

Type D



15



Uncertain

Code

- Fiber tail
- Fiber shaft
- Fiber knob
- New trimerisation motif
- Linker motif
- EGF
- Native trimerisation motif

- (Gly4Ser)4 linker
- Turn b from repeat 17 of Ad5 fiber shaft
- Turn b from repeat 22 of Ad5 fiber shaft

Fig. 4b

664720-2294103

THIS PAGE BLANK (USPTO)